

FIFRA 2(ee) Recommendation for the Control of American Beech (*Fagus grandifolia*) Within Forestry Sites

ACCEPTED

Oct 22, 2012

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration

Rodeo

EPA Reg. No. 62719-324

Classified for
“RESTRICTED USE”
in New York State
under 6NYCRR Part 326

Doc id: 533204

Directions for Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of pesticide application.
- For cut stump application, apply Rodeo using a 50 to 100 percent solution to freshly cut surface immediately after cutting. Apply product using suitable equipment to ensure coverage of the entire cambium.
- For injection and frill applications, apply a 25 to 100 percent solution to a continuous frill or evenly spaced cuts around the tree below all branches. Apply using suitable equipment which must penetrate into living tissue. Avoid application techniques that allow runoff to occur from frill or cut areas.
- For foliar applications, use the application rates consistent with the application method listed on the label. In forestry site applications, a non-ionic surfactant containing greater than 80 percent active ingredient is required for foliar applications of Rodeo.
- Rodeo may be tank mixed with Oust, Arsenal, and Garlon products according to label instructions for foliar applications within forestry sites.
- For best results applications should be made during periods of active growth, after full leaf expansion and before fall color and leaf drop.
- The use of colorants, dyes, and/or surfactants must comply with the information on the product label.
- Follow all applicable directions, restrictions, and precautions on the product label.
- Do not apply more than 8 quarts of Rodeo per acre per year.